AFRL-SR-BL-TR-02-

0050

REPORT DOCUMENTATION PAGE

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instruction needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any chier aspect of the burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jeffers on Davis Highway, Suite 12, Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank) 2 PEROFT DATE

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE	3. REPORT TYPE AND	
4. TITLE AND SUBTITLE	January 14, 2002	Final Technical Rep	ort 1 Jun 97-31 Dec 00 5. FUNDING NUMBERS
VERY LOW FREQUENCY (V	LF) REMOTE SENSING OF G BY THE AURORAL ELECTRO		G F49620-97-1-0468
6. AUTHOR(S)			1
T. F. Bell, U. S. Inan, M. K. Demirkol, M. W. Chevalier, and R. Moore			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			8. PERFORMING ORGANIZATION REPORT NUMBER
STAR Laboratory, Packard Buildi Stanford University	ing, Stanford, California	94305	None
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING / MONITORING
			AGENCY REPORT NUMBER
Department of the Air Force AFOSR/NE	110 Duncan Avenue Room B115		
AFOSIONE	Bolling AFB		
	DC 20332-8050		
11. SUPPLEMENTARY NOTES			
		AIR FORC	PEOFFICE OF SCIENTIFIC PEOFAPORISM CODE THIS TECHNICAL TEC
12a. DISTRIBUTION / AVAILABILITY STATEMENT HAS BEEN			DESCRIPTION OF THE TRANSPORT OF THE PROPERTY O
DISTRIBUTION STATEMENT A LAWAFR 100			REVIEWED AND IS A PPROVED FOR FUBLIC RELEASE 90-12. DISTRIBUTION IS UNLIMITED.
Approved for Public Release			SU-12. DISTRIBUTION IS LIN MATER
modified D-region when perturbed sub-ionospheric wave phase and a the North American continent. A t captured that exhibited features co amplitudes were pertured by as mu	the VLF remote diagnostic technical by atmospheric gravity waves graphitude measurements at four stotal of ten separate propagation ensistent with impulsive generation uch as 3 dB by the gravity wavement of the D-region driven by the state of the design of the property wavement of the property wavement of the property wavement of the property driven by the state of the property wavement of the	generated by the aurora sites located in Alaska paths were monitored. on by the auroral electral s. The measured VLF whe gravity waves. We	der to characterize both the ambient and l electrojet. The characterization involved VLF and four sites located along the east coast of A large number of gravity wave events were rojet. Measured VLF sub-ionosperic wave wave amplitude and phase perturbations were conclude that gravity waves excited by the oheric VLF waves.
	20	020221	055
14. SUBJECT TERMS			15. NUMBER OF PAGES
Gravity waves, sub-ionospheric VLF waves			16. PRICE CODE
17. SECURITY CLASSIFICATION	18. SECURITY CLASSIFICATION	19. SECURITY CLASSI	FICATION 20. LIMITATION OF ABSTRACT
OF REPORT UNCLASSIFIED	OF THIS PAGE UNCLASSIFIED	OF ABSTRACT UNCLASSI	

UNCLASSIFIED

FINAL TECHNICAL REPORT

SPONSOR: DEPARTMENT OF THE AIR FORCE

SPONSER REF: F49620-97-1-0468

TITLE: VERY LOW FREQUENCY (VLF) REMOTE SENSING OF GRAVITY WAVES GENERATED BY THE AURORAL ELECTROJET

PERIOD OF PERFORMANCE: 06/01/97 - 12/31/00

PRINCIPLE INVESTIGATOR: PROFESSOR U. S. INAN

STAR LABORATORY, STANFORD UNIVERSITY

Report prepared by: Timothy F. Bell Senior Research Associate STAR Laboratory, Stanford University January 14, 2002

SUMMARY

During the period of performance the VLF remote diagnostic technique was applied in order to characterize both the ambient and modified D-region when perturbed by atmospheric gravity waves generated by the auroral electrojet. The characterization involved VLF sub-ionospheric wave phase and amplitude measurements at four sites located in Alaska and four sites located along the east coast of the North American continent. A total of ten separate propagation paths were monitored. A large number of gravity wave events were captured that exhibited features consistent with impulsive generation by the auroral electrojet. Measured VLF sub-ionosperic wave amplitudes were pertured by as much as 3 dB by the gravity waves. The measured VLF wave amplitude and phase perturbations were consistent with a vertical displacement of the D-region driven by the gravity waves. We conclude that gravity waves excited by the auroral electrojet can have significant effects upon sub-ionospheric VLF signal amplitude and phase.

FINAL TECHNICAL REPORT

1. Contract Purpose

The purpose of this contract (F49620-97-1-0468) was to carry out a program of technique development, data acquistion, and data analysis in order to characterize both the ambient and modified D-region when perturbed by atmospheric gravity waves generated by the auroral electrojet. The data was acquired at eight sites which were established and supported by the Parent contract F19628-96-C-0149-P00003. The data acquired at these sites consisted of measurements of the phase and amplitude of sub-ionospheric VLF signals transmitted by VLF communications transmitters operated by the United States Navy.

2. Period of Performance

The original period of performance under this contract extended from June 1, 1997, through May 31, 2000. However, in order to better fit the academic schedule, a no cost seven month extension of the Grant was requested and granted. Thus the actual period of performance was June 1, 1997, through December 31, 2000.

3. Work Provided

During the period of performance of this grant, Stanford University acquired sub-ionospheric VLF wave amplitude and phase data during atmospheric gravity wave events at each of the Alaskan and East Coast sites supported by the Parent contract. Instrumentation used for this task was developed and deployed under the sponsorship of the Parent contract F19628-96-C-0149-P00003. The gravity wave data was analyzed using general algorithms developed under the Parent contract and specific algorithms developed for gravity wave analysis. This work was reported in a paper delivered at the December American Geophysical Union in 1998 [Demerkol, et al., 1998]. The complete citation for this paper is given in Section 4.

4. Results of atmospheric gravity wave study

As mentioned above, results of the atmospheric gravity wave study were reported in a paper delivered at the December American Geophysical Union in 1998 [Demirkol, M. K., U. S. Inan, and T. F. Bell, Quantitative determination of D-region density and temperature profiles via inversion of sub-ionospheric VLF measurements, *EOS*, 79, F674, 1998]. A number of typical examples of gravity wave events were observed on November 2000 at the Delta Junction VLF receiving site. Some of these events involved the subionospheric signal from the NPM transmitter. Figure 1 shows the path of the VLF signal from the NPM tranmitter in Hawaii as it propagated to the VLF receiving site at Delta Junction in Alaska. Delta Junction is located at a position close to the auroral electrojet where gravity waves are often generated.




Fig. 1. Propagation path of the NPM VLF signal from Hawaii across Alaska to the Delta Junction (DJ) VLF receiving site. The footprints of the magnetic shells L=4 and 7 are indicated by dashed lines.

Figure 2 shows the amplitude and phase of the sub-ionospheric NPM signal as observed at

Delta Junction during the period: 0510 UT to 0710 UT. Quasi-sinusoidal oscillations in both amplitude and phase are observed as a gravity wave moves across the NPM propagation path. The largest amplitude change due to the gravity wave is approximately 3 dB. The maximum phase change is approximately 15 degrees.




Fig. 2. The upper and lower panels show respectively the amplitude (in normalized units) and the phase of the NPM sub-ionospheric VLF signal as observed at Delta Junction.

5. List of personnel contributing to report

Tim Bell

Mike Chevalier

Umran Inan

Mike Johnson

Robb Moore